

Growth Inhibitory Activity of *Clinopodium Nepeta* Essential Oil against Gram-positive and Gram-negative Bacteria: *in vitro* Studies

To Cite:

Chouitah O, Mandal S. Growth Inhibitory Activity of *Clinopodium Nepeta* Essential Oil against Gram-positive and Gram-negative Bacteria: *in vitro* Studies. *Discovery*, 2022, 58(321), 1041-1044

Author Affiliation:

¹BGSS Laboratory of Engineering and Health Security, University de Mascara- 29000, Algeria (Centre National De Recherche Scientifique Et Technique, Rabat, Maroc): E-mail: chouitah_o@yahoo.fr

²Laboratory of Microbiology and Experimental Medicine, Department of Zoology, University of Gour Banga, Malda-732103, India

***Correspondence:**

Dr. Shyamapada Mandal, Professor, Department of Zoology, University of Gour Banga, Malda-732103, India. E-mail address: sam.micro11@ugb.ac.in ; samtropmed@gmail.com

Peer-Review History

Received: 23 July 2022

Reviewed & Revised: 26/July/2022 to 24/August/2022

Accepted: 27 August 2022

Published: September 2022

Peer-Review Model

External peer-review was done through double-blind method.



© The Author(s) 2022. Open Access. This article is licensed under a Creative Commons Attribution License 4.0 (CC BY 4.0), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Ourida Chouitah¹, Shyamapada Mandal^{2*}

ABSTRACT

Present research reports chemical composition and antibacterial activity essential oil from aerial parts of *Clinopodium nepeta* growing widely in Mascara, Algeria. It is an ornamental plant with many medicinal usages. Antimicrobial activity of essential oil extracted by hydro-distillation process from *Clinopodium nepeta* was tested on gram-negative (*Escherichia coli*, *Pseudomonas aeruginosa*) and gram-positive (*Staphylococcus aureus* and *Bacillus subtilis*) bacteria. The essential oil yield from *Clinopodium nepeta* was 0.54–0.68% (v/w), which exhibited antibacterial activity against the test bacteria displaying zone diameter of inhibition ranging from 14 – 17 mm. Thus, the plant might be an important source of active ingredients useful to combat bacterial infection to humans.

Keywords: *Clinopodium nepeta*, essential oil, antibacterial activity, gram-negative bacteria, gram-positive bacteria

1. INTRODUCTION

Medicinal and aromatic plants are used for treatment as alternative therapies (Blowman et al., 2018; Das and Mandal, 2022). Plant essential oils are useful to detoxify and to promote the human health. They are also excellent sources of natural preservatives with antioxidant potential and promising antimicrobial agents against bacterial pathogens (Mandal and Mandal, 2016). Their chemical composition includes primarily terpenoids, like monoterpenes, sesquiterpenes, diterpenes, acids, alcohols, aldehydes (Masango, 2005).

In the last few years, a worrisome increase in antibiotic resistance of pathogenic bacteria has been observed (Levy, 2000). The Bacterial resistance is a global health problem with significant impact on mortality rates and health care costs (Boucher et al., 2009). Evidence has indicated that multi-drug resistant (MDR) bacterial strains can become increasingly virulent, hindering the treatment of several infections (Mandal, 2015). Resistances among gram-positive and gram-negative pathogens that cause infection in the hospital and in the community

have been reported, and bacteria such as *Staphylococcus aureus*, *Escherichia coli*, and *Pseudomonas aeruginosa* have been recognized as highly pathogenic microorganisms responsible for a significant number of serious infections (Mogasale et al. 2021). Based upon the emergence of antibiotic resistant bacteria, priority in the search for new antimicrobial agents has been given for combating MDR pathogenic bacteria (Demirci et al., 2011).

In the folk medicine of different countries of the world, *Clinopodium nepeta* L. (Lamiaceae family), belonging to the mint family: Lamiaceae (with 236 genera is a large dicotyledonous group with a high number of aromatic species), has been widely used for the treatment of insomnia, depression, respiratory and gastroenteric diseases (Saltos et al., 2014). Its essential oil is extensively employed to add a distinctive aroma and flavour to food and can be used in cosmetics or pharmaceuticals for an enormous number of ailments (Chouhan et al., 2017).

The genus *Clinopodium* consist of flowering plants, widely distributed in southern and southeastern Europe, North America, Latin America and Asia. This aromatic plant, reaching 80 cm in height (Dorman et al., 2000). It has the appearance of a small shrub of lilac flowers, with small and ovals leaves and it is widely distributed in Mediterranean region (Bozovic 2017). The earlier authors have also reported the antibacterial activity of *Clinopodium nepeta* essential oil (CNEO) against both gram-negative and gram-positive bacteria, and demonstrated higher activity of the essential oil against *Bacillus cereus* and *Streptococcus sanguinis* (Ozturk et al., 2021). The present study determines the antibacterial activity of CNEO as a non-traditional raw material in pharmaceuticals for the treatment of bacterial infection.

2. MATERIAL AND METHODS

The aerial parts including leaves of *Clinopodium nepeta* were collected in Ain Fares Mascara region, and the Voucher specimens was identified and deposited at the herbarium of aromatic plants of the University of Mascara, Algeria, with the number 5620. Plant materials were dried at room temperature (20–25 °C), and were grinded. Essential oil was obtained by hydro-distillation method (Debbabi et al., 2022), using air-dried ground plant materials of *Clinopodium nepeta* using 50 g of dry leaves in 1 L of distilled water and extracted for 3h.

The essential oil was dried over anhydrous sodium sulfate and stored in a scaled vial in the dark at 4°C (Adams, 2007). The yield of CNEO was calculated using a dry weight gravimetric method. Some parameters (point of freezing, refractive index, solubility in ethanol at 90 °C, and the acidity) of CNEO were measured.

The antibacterial activity of the CNEO was determined following the method, as described earlier (Mandal et al., 2007; Sircar and Mandal (2016), using sterile paper discs (6 mm diameter) impregnated with CNEO. The bacterial strains used in the study were *Escherichia coli* ATCC 3560 (from urine), *Bacillus subtilis* ATCC 4350 (from soil), *Pseudomonas aeruginosa* ATCC 27853 (from Nasal swab), and *Staphylococcus aureus* ATCC 14990 (from rectal swab).

3. RESULT AND DISCUSSION

The hydro-distillation of *Clinopodium nepeta* produced a light-yellow color essential oil. The yield percentage of *Clinopodium nepeta* aerial parts essential oil was 0.13%, with characteristic flavour, and the leaf essential oil yields up to 0.54–0.68% (v/w). Determination of density was done by double weighing $d = 0.865 \text{ g/cm}^3$, the optical activity N20 by polarimetry was +10.12, the refractive index $n = 1.4449$ by an interferometric method. Ester index was 0.78 and freezing point of essential oil was -19°C (Table 1). The CNEO demonstrated bactericidal activity against both gram-positive (*Staphylococcus aureus* and *Bacillus subtilis*) and gram-negative (*Escherichia coli* and *Pseudomonas aeruginosa*) bacterial strains (Figure 1).

Table 1. Physicochemical property of *Clinopodium nepeta* essential oil

• Specification	Parametric value
• Density D20	0.865 g/cm ³
• Refractive index	1.4449
• Optical activity N20	+10.12
• Solubility in ethanol (90 %)	1 :1
• Freezing point (°C)	-19
• Ester index	0.78




Figure 1. Antibacterial activity in terms of ZDI (zone diameter of inhibition) of *Clinopodium nepeta* essential oil against gram-negative and gram-positive bacteria

The essential oil had a strong antimicrobial activity against gram-positive as well as gram-negative bacteria (D'Aquila et al., 2022). As per the earlier report, the antibacterial susceptibility of CNEO as was determined by disc diffusion, the all microorganisms tested were sensitive to essential oil (Kitic et al., 2002). Essential oil exhibited wide-ranging inhibitory activities against various microbial pathogens and other organisms (Debbabi et al., 2022), with the penetration of lipid layer of the bacterial cell membrane thereby disrupting the bacterial cell wall structure. The various bioactive components present in the CNEO might damage the cell membrane and decrease bacterial metabolic activity (Kitic et al., 2002; Marongiu et al., 2010; Ozturk et al., 2021). As demonstrated by Kerbouche et al. (2013), *Staphylococcus aureus* had highest sensitivity with ZDI of 40 mm, while *Pseudomonas aeruginosa* had lowest susceptibility to the test essential oil from *Satureja calamintha* subsp. *Nepeta*. The current study provides the benefit of using CNEO as a natural food preservative by inhibiting growth of potential food borne pathogenic, such as *Staphylococcus aureus* (ZDI: 14 mm) and *Bacillus subtilis* (ZDI: 16 mm), *Escherichia coli* (ZDI: 14 mm) and *Pseudomonas aeruginosa* (ZDI: 17 mm). Thus, CNEO showed antibacterial activity against both gram-negative and gram-positive bacteria, with highest inhibition against *Pseudomonas aeruginosa*.

4. CONCLUSION

The antibacterial potentiality of *Clinopodium nepeta* essential oil suggests its capacity of usefulness in pharmaceuticals for the preparation of biotherapeutics active against pathogenic bacteria. However, analysis for bioactive compounds along with their pharmacological properties and mechanism of action are warranted, for which further research is ongoing.

Funding and Acknowledgement

The first author (OC) of this article acknowledges the support of the Ministry of High School and Scientific Research of the Republic of Algeria, PRFU project: D01NO1UN290120190001, for the current research.

Conflicts of interests

The authors declare that there are no conflicts of interests.

Data and materials availability

All data associated with this study are present in the paper.

REFERENCES AND NOTES

1. Adams RP. (2007). Identification of essential oil components by gas chromatography/mass spectrometry, 4th Edition.

- Allured Publ., Carol Stream, IL.
2. Blowman K, M. Magalhães, M. F. L. Lemos, C. Cabral, I. M. Pires (2018). Anticancer properties of essential oils and other natural products. Evidence-Based Complementary and Alternative Medicine. doi: 10.1155/2018/3149362
 3. Boucher HW, Talbot GH, Bradley JS, et al. (2009). Bad bugs, no drugs: no ESCAPE! An update from the infectious diseases society of America. Clin Infect Dis, 48, 1-12.
 4. Chouhan S, Sharma K, Guleria S. Antimicrobial activity of some essential oils—present status and future perspectives. Medicines 2017, 4, 58; doi:10.3390/medicines4030058
 5. D'Aquila P, Paparazzo E, Crudo M., Bonacci S, Procopio A, Passarino G, Bellizzi D (2022). Antibacterial activity and epigenetic remodeling of essential oils from Calabrian aromatic plants. Nutrients, 14, 391. doi: 10.3390/nu14020391
 6. Das RC, Mandal S (2022). Antibacterial activity of some plant extracts alone and in combination with antibiotics against multidrug resistant potential pathogenic bacteria from sewage water in Kolkata, India. Discovery, 58(318), 579-587.
 7. Debbabi H, El Mokni R, Chaieb I, Nardoni S, Maggi F, Caprioli G, Hammami S (2020). Chemical Composition, Antifungal and Insecticidal Activities of the Essential Oils from Tunisian *Clinopodium nepeta* subsp. *nepeta* and *Clinopodium nepeta* subsp. *glandulosum*. Molecules 25(9): 2137. doi: 10.3390/molecules25092137
 8. Demirci B, Temel HE, Portakal T, et al. (2011). Inhibitory effect of *Calaminthana nepeta* subsp. *glandulosa* essential oil on lipoxygenase. Turk. J. Biochem., 36, 290-295.
 9. Dorman HJD, Deans SG (2000). Antimicrobial agents from plants: Antibacterial activity of plant volatile oils. *J. Appl. Microbiol.*, 88, 308-316.
 10. Kerbouche L., Hazzit M., Baaliouamer A (2013). Essential oil of *Satureja calamintha* subsp. *nepeta* (L.) Briq. from Algeria: Analysis, antimicrobial and antioxidant activities. J. Biol. Act. Prod. Nat., 3: 266-272.
 11. Kitic, D., Jovanovic, T., Ristic, M., Palic, R., & Stojanovic, G. (2002). Chemical composition and antimicrobial activity of the essential oil of *Calamintha nepeta* (L.) Savi ssp. *glandulosa* (Req.) P. W. Ball from Montenegro. Journal of Essential Oil Research, 14, 150-152.
 12. Levy, S.B. The 2000 Garrod Lecture. Factors impacting on the problem of antibiotic resistance. J. Antimicrob. Chemother 2002. 49: 25-30.
 13. Mandal M, Mandal S, Pal NK. (2007). Antibacterial potential of *Azadirachta indica* seed and *Bacopa monniera* leaf extracts against multidrug resistant *Salmonella enterica* serovar Typhi isolates. Arch Med Sci 3, 14-18.
 14. Mandal M, Mandal S. (2016). Cumin (*Cuminum cyminum* L.) Oils. In: Essential Oils in Food Preservation, Flavour and Safety, 377-382. Elsevier, London. doi: 10.1016/B978-0-12-416641-7.00042-0
 15. Mandal S. (2015). Can over-the-counter antibiotics coerce people for self-medication with antibiotics? Asian Pacific Journal of Tropical Disease, 5, S184-S186. doi:10.1016/S2222-1808(15)60886-6
 16. Masango P. (2005). Cleaner production of essential oils by steam distillation. J. Clean. Prod, 13: 833-839.
 17. Mogasale VV, Saldanha P, Pai V, Rekha PD, Mogasale V (2021). A descriptive analysis of antimicrobial resistance patterns of WHO priority pathogens isolated in children from a tertiary care hospital in India. Scientific Reports, 11: 5116.
 18. Saltos MBV, Puente BFN, Malafronte N, Braca A (2014). Phenolic Compounds from *Clinopodium tomentosum* (Kunth) Govaerts (Lamiaceae). J. Braz. Chem. Soc., 25, 11, 2121-2124.
 19. Sircar B, Mandal S. (2016). Antibacterial Activity of *Mimusops elengi* Leaf, Seed and Bark Extracts Alone and in Combination with Antibiotics against Human Pathogenic Bacteria. Transl Med 6: 187. doi:10.4172/2161-1025.1000187